

II. Combining events: Multiplication and addition

Unions of events

- $A \cup B$ is the set of outcomes in A or B , meaning at least one of A or B is true.
- This is the inclusive OR: It means one or the other or both. (The exclusive OR, meaning one or the other but not both, is not as common.)

Think of soup or salad at a restaurant. This isn't that.

Draw a Venn diagram with S , A , and B .

- $$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$
- If A and B are disjoint events (no overlap), then $P(A \cap B) = 0$, so we get

$$P(A \cup B) = P(A) + P(B)$$

Draw a Venn diagram with S , A , and B .

Intersections of events

- $A \cap B$ is the set of outcomes in A and B , meaning both of A and B are true.

Draw a Venn diagram with S , A , and B .

- To find $P(A \cap B)$, the multiplication rule is often useful:
- If you have m possible outcomes for one experiment and n possible outcomes for a second, independent experiment, then there are mn possible combined outcomes.

Conditional probability

- The conditional probability $P(A|B)$ (“ $P(A$ given B)”) is the probability of event A , given that event B is true.

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Draw a Venn diagram with S , A , and B .

Independence

- Events A and B are independent if knowing that B is true would not change the probability that A is true:

$$P(A) = P(A|B)$$

- **Warning:** Independent does not mean disjoint. (Disjoint means that $P(A \cap B) = 0$.)
- If A and B are independent, then $P(A) = \frac{P(A \cap B)}{P(B)}$, so we get

$$\boxed{P(A \cap B) = P(A)P(B)}.$$

Example I

Choose a whole number at random from 1 to 100. What is the probability that it will be even or greater than 80?

By counting:

$$\begin{aligned} S &:= \{\text{all numbers}\} \\ A &:= \{\text{even numbers}\} \\ B &:= \{\text{numbers} > 80\} \\ |A| &= 50 \\ |B| &= 20 \\ |A \cap B| &= 10 \\ |A \cup B| &= |A| + |B| - |A \cap B| = 50 + 20 - 10 = 60 \\ P(A \cup B) &= \frac{|A \cup B|}{|S|} = \frac{60}{100} = \boxed{\frac{3}{5}} \end{aligned}$$

By probability:

$$\begin{aligned} P(A) &= \frac{50}{100} = \frac{1}{2} \\ P(B) &= \frac{20}{100} = \frac{1}{5} \\ P(A \cap B) &= \frac{10}{100} = \frac{1}{10} \\ P(A \cup B) &= P(A) + P(B) - P(A \cap B) = \frac{1}{2} + \frac{1}{5} - \frac{1}{10} = \frac{5 + 2 - 1}{10} = \boxed{\frac{3}{5}} \end{aligned}$$

Example II

A combination deal at a restaurant includes a main dish, drink, and dessert. If the restaurant serves four types of drinks, 12 main dishes, and three desserts, then how many different combination meals are possible?

$$4 \cdot 12 \cdot 3 = \boxed{144}$$

Example III

We roll two dice, a red one and a blue one. What is the probability that the red one shows a 4 and the blue one is odd?

By counting:

$$P(A \cap B) = \frac{|A \cap B|}{|S|} = \frac{|\{41, 43, 45\}|}{36} = \boxed{\frac{1}{12}}$$

By probability:

These are independent, since knowing that one is true would not change the the probability that the other is true.

$$P(A \cap B) = \left(\frac{1}{6}\right) \left(\frac{1}{2}\right) = \boxed{\frac{1}{12}}$$

Example IV

Flip a coin four times. What is the probability that there will be at least three heads, given that the first two flips are heads?

$$S := \{\text{all 16 outcomes}\}$$

$$A := \{\text{at least three heads: HHHT, HHTH, HTHH, THHH, HHHH}\}$$

$$B := \{\text{first two are heads: HHHH, HHHT, HHTH, HHTT}\}$$

$$A \cap B = \{HHHT, HHTH, HHHH\}$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{3}{16}}{\frac{4}{16}} = \boxed{\frac{3}{4}}$$

Example V

Roll two dice (red and blue). Define the events:

A = red die is 3

B = total is 7

C = total is 8

Calculate the conditional probabilities:

$P(B|A)$

$P(A|B)$

$P(C|A)$

$P(A|C)$

$P(C|B)$

$P(B|C)$

$$\begin{aligned}
 A = \text{red die is 3.} \quad P(A) &= \frac{6}{36} = \frac{1}{6}. \\
 B = \text{total is 7.} \quad P(B) &= \frac{6}{36} = \frac{1}{6}. \\
 C = \text{total is 8.} \quad P(C) &= \frac{5}{36}.
 \end{aligned}$$

$$P(B|A) = \boxed{\frac{1}{6}}$$

$$P(A|B) = \boxed{\frac{1}{6}}$$

$$P(C|A) = \boxed{\frac{1}{6}}$$

$$P(A|C) = \boxed{\frac{1}{5}} \quad \text{Different from } P(C|A)!$$

$$P(C|B) = \boxed{0}$$

$$P(B|C) = \boxed{0}$$

Example VI

As above, roll two dice (red and blue). Define the events:

$$A = \text{red die is 3}$$

$$B = \text{total is 7}$$

$$C = \text{total is 8}$$

Are events A and B independent? A and C ? B and C ?

By the formulas, A and B are independent. A and C are dependent. B and C are dependent.